

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/053623

International filing date: 21 December 2004 (21.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: EP  
Number: 03104954.7  
Filing date: 23 December 2003 (23.12.2003)

Date of receipt at the International Bureau: 04 February 2005 (04.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**


03104954.7

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**





Anmeldung Nr.:  
Application no.: 03104954.7  
Demande no:

Anmeldetag:  
Date of filing: 23.12.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Akzo Nobel N.V.  
Velperweg 76  
6824 BM Arnhem  
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

COMBINATION VACCINE FOR POULTRY

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A61K39/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI



## Combination vaccine for poultry

The present invention relates to a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*, to the use of a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus for the manufacturing of such a combination vaccine, to methods for the preparation of said combination vaccine and to vaccination kits for the immunization of poultry against *Ornithobacterium rhinotracheale*.

Over the last decades, in many countries a strong raise in both the number of poultry farms, and in addition, an increasing number of animals per farm has been seen. This situation has a serious consequence: large scale outbreaks of poultry diseases are seen more and more frequently. This in turn has caused an increasing need for new and better vaccines and vaccination programs in these countries.

Nowadays, most animals are immunized against a number of diseases of viral, bacterial and parasitic origin. Examples of bacterial agents infective to poultry are *Ornithobacterium rhinotracheale*, *Haemophilus paragallinarum* (Coryza), *Salmonella spp*, *Pasteurella multocida*, *Bordetella bronchiseptica* and *E. coli*. Examples of viral poultry pathogens are Newcastle Disease virus (NDV), Infectious Bronchitis virus (IBV), Turkey Rhinotracheitis virus (TRT), Herpesvirus of Turkeys (HVT), Fowlpox virus (FPV), Avian Reovirus (ARV), Infectious Laryngotracheitis virus (ILT), Marek's Disease Virus (MDV) and Infectious Bursal Disease virus (IBDV).

*Ornithobacterium rhinotracheale* is a relatively new bacterium causing a disease known for about a decade now, and found frequently in i.a. chickens and turkeys. Clinical signs in chickens are e.g. airsacculitis or coughing, pneumonic lungs or pleuritis. In turkey flocks in several parts of the world, a comparable infection of the respiratory tract is found. Mortality in flocks suffering from the disease can be as high as 5%. The first clinical signs are comparable to infection in chicken: sneezing and nasal discharge. In some animals clinical signs of acute infection are seen. Examination of sacrificed animals shows edema of the lungs, fibrinopurulent pneumonia and often serofibrinous pericarditis and serofibrinous infection of the airsacs. *Ornithobacterium rhinotracheale* is extensively described in European Patent EP0.625.190. Identification, serotyping and experimental infection in turkeys and chickens have been described e.g. by van Empel, P.C.M. et al., in Journ. of Clin. Microbiol. 35: 418-421 (1997), by van Empel, P.C.M. et al., in Avian Diseases 40: 858-864 (1996) and by van Empel, P.C.M. et al., in Avian Pathology 28:217-227 (1999). A review on *Ornithobacterium rhinotracheale* has been published in Avian Pathology 28: 217-227 (1999) by van Empel, P.C.M. and Hafez, H.M.

If an animal suffers from infection with a virulent pathogen, the immune system will try to remove the pathogen. Provided that the infected animal survives the infection, it usually develops a long-lasting immunity against the pathogen.

- 5 Vaccination with an attenuated pathogen mimics the natural infection in that it induces immunity, however without causing unacceptable clinical signs. Some clinical signs, in other words; some residual virulence, has to be accepted however because without this, the immune system will in most cases be insufficiently triggered. Generally spoken, vaccination is the way to avoid large scale infection and its negative consequences.

10

Vaccines against most of the known pathogenic bacteria and pathogenic viruses may comprise either inactivated or live attenuated pathogens, depending upon the purpose and the way of administration. But if it comes to efficient and early protection of young animals, combined with ease of application, live attenuated vaccines are clearly an attractive choice.

- 15 Such vaccines can be given from the day of birth, for poultry even *in ovo*, and if preferred, they can be applied from day of birth simply by spraying or through drinking-water application.

20

Spraying of vaccines is especially attractive for poultry and is currently applied on a very large scale for e.g.: Infectious Bronchitis virus, Newcastle Disease virus, Turkey Rhinotracheitis virus, Infectious Bursal Disease virus and Avian Reovirus.

The main disadvantage of vaccines derived from virulent pathogens is, that they must be carefully and sufficiently attenuated in order to be safe.

- 25 This requirement can relatively easy be fulfilled for highly virulent microorganisms, because they can in most cases be highly attenuated while remaining their vaccinating (immunity-inducing) capacities.

30

Of the bacteria mentioned above, *Ornithobacterium rhinotracheale* belongs to the so-called secondary pathogens. A secondary pathogenic bacterium is a bacterium that does not readily cause disease in healthy animals. As a consequence, the immune system will not or insufficiently be triggered and therefore, no immunity will develop.

In animals kept under sub-optimal conditions such as high stock density and stress, such secondary pathogenic bacteria can however cause dramatic morbidity and mortality.

35

Thus, for secondary pathogens, the design of a live attenuated vaccine is severely hampered by the unpredictable behavior of the bacteria. Unpredictable in the sense that the level of virulence of such bacteria does not solely depend on their level of attenuation, but (contrary to

what is the case for primary pathogens) also largely on the physical situation of the animal to be vaccinated.

This is a real dilemma as will be sketched below using *Ornithobacterium rhinotracheale* as example.

5

*Ornithobacterium rhinotracheale* is a relatively mild bacterium, in the sense that in perfectly healthy Specified Pathogen Free (SPF) poultry kept under optimal conditions it might be difficult to induce disease at all, even when using non-attenuated, wild-type, bacteria.

Thus, such poultry, be it chickens, turkeys or ducks, even when "vaccinated" with the wild-type bacterium, are not affected and as a consequence they do not build up any protection against *Ornithobacterium rhinotracheale*. An example of a wild-type strain is the *Ornithobacterium rhinotracheale* strain 3263/91 as deposited at the Centraalbureau voor Schimmelcultures (CBS), Oosterstraat 1, PO.box 273, 3740 AG Baarn, The Netherlands, under accession-number 400.92.

15

However, poultry kept under standard conditions for commercial breeding, i.e. a high stock density, stress, poor ventilation and/or high ammonia levels, are much more vulnerable to *Ornithobacterium rhinotracheale*. They are severely affected by wild-type *Ornithobacterium rhinotracheale* strains.

Especially if such animals would already suffer from another infection the effect could be dramatic. In animals suffering from (sometimes even sub-clinical) infections with other bacteria or viruses, the wild-type bacterium often causes massive and severe morbidity.

25

In addition to this and unlike other secondary pathogenic bacteria, the pathological effects of any *Ornithobacterium rhinotracheale* strain in poultry not only depend on the virulence of the strain as such and the physical condition of the infected animal but, especially in the case of chickens, also heavily upon the chicken species. It is e.g. well-known in the art, that broilers are much more susceptible to *Ornithobacterium rhinotracheale* infection than layers.

30

For the reasons given above, the development of a live attenuated vaccine starting from a secondary pathogen in general and especially *Ornithobacterium rhinotracheale*, although possible, is complex to say the least. On the one hand, even a wild-type strain of these bacteria is not capable of inducing protection in perfectly healthy, less susceptible animals, whereas on the other hand even an attenuated strain is only suitable in some cases for use in susceptible animals or animals suffering from stress and other negative environmental factors, the suitability highly depending upon the level of stress and the poultry species.

35



Only a live over-attenuated *Ornithobacterium rhinotracheale* strain would be safe under all circumstances in poultry industry and in both broiler chickens and layer chickens.

5 An attenuated strain, let alone an over-attenuated strain, will however not trigger any immune reaction in healthy, less susceptible animals, because in such animals even the virulent strain does not trigger an immune reaction.

10 This dilemma makes it difficult to provide vaccines based upon live secondary pathogens, more specifically *Ornithobacterium rhinotracheale*, that are both efficacious and safe, regardless the kind of animal to be vaccinated and its physical condition.

It is an objective of the present invention to provide vaccines that add to solving this dilemma.

15 In order to reach this goal, the present invention provides a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale* wherein the combination vaccine comprises a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus.

20 The present invention relates to the surprising finding that even over-attenuated secondary pathogenic *Ornithobacterium rhinotracheale* bacteria are well capable of inducing a protective response provided that they are given in combination with a live attenuated poultry virus. Such over-attenuated secondary pathogenic bacteria are inherently safe because they only cause moderate clinical signs even when they would accidentally be administered at a moment in time at which a field infection with a wild-type virus has occurred.

25 An over-attenuated *Ornithobacterium rhinotracheale* bacterium is a bacterium that (contrary to an attenuated *Ornithobacterium rhinotracheale* strain) is not capable of inducing a protective immune response to *Ornithobacterium rhinotracheale* in primed animals. A primed animal is an animal that has received, prior to administration of the vaccine, a virulent pathogen that predisposes for infection with a secondary pathogenic bacterium.

30 Primed animals provide an objective animal model for testing the level of attenuation. Animals randomly taken from commercial farms would not provide a stable model, because inherently, their physical condition is difficult to establish and highly variable. Primed animals, contrary to this, are of known SPF-origin, they are by definition in good health and they received, as a priming, a well-defined amount of a well-defined pathogen.

35 The priming virus of choice for the chicken model is Newcastle Disease virus. Turkey Rhinotracheitis virus is the priming virus of choice in turkeys. Thus, in order to see if an attenuated *Ornithobacterium rhinotracheale* bacterium for use in chickens indeed behaves

over-attenuated in chickens, this should be tested in chickens primed with Newcastle disease virus, as is e.g. indicated in the Examples section.

A protective immune response is an immune response that gives, after challenge, a

- 5 statistically significant decrease in respiratory tract lesion score of equal or more than 50% compared to non-vaccinated animals. The respiratory tract lesion score test is explained in the Examples section.

The nature of the mutation(s) leading to the over-attenuated behavior is not critical. Many

- 10 attenuating mutations known in the art are suitable. Suitable *Ornithobacterium rhinotracheale* mutations are e.g. the classical PurD-, RecA- and Aro-mutations, known in the art for many bacterial species. Also, many temperature-sensitive mutants are suitable. Of these, PurD- and RecA-mutations are preferred. The PurD gene encodes a protein that is involved in the Purine ribonucleotide biosynthesis. A PurD-mutation is defined as a mutation that disturbs the Purine ribonucleotide biosynthesis in the sense that such a mutant makes less or less active or no
- 15 PurD gene product compared to wild type *Ornithobacterium rhinotracheale* strains.

The RecA gene encodes a protein that is involved in recombination processes. A RecA-mutation is defined as a mutation that disturbs recombination processes in the sense that such a mutant makes less or less active or no RecA gene product compared to wild type *Ornithobacterium rhinotracheale* strains.

- 20 Mutations such as PurD- and RecA-mutations are most easily obtained by simply deleting the gene encoding PurD and RecA. Such (site-directed) mutation techniques are well-known in the art.

The level of attenuation of the live attenuated viral component of the combination vaccine according to the invention is not critical. Usually, a live attenuated viral vaccine strain will be chosen that is suitable as a vaccine, i.e. a strain that induces protection against infection without causing an unacceptable level of pathogenicity. Many live attenuated viral poultry vaccines are commercially available. Such strains can simply be purchased from various producers.

30

Suitable live attenuated poultry viruses are e.g. Infectious Bronchitis virus, Turkey Rhinotracheitis virus, Newcastle Disease virus, Avian Reovirus and Marek's virus.

- 35 Examples of suitable live attenuated viral vaccine strains especially suitable in chickens are e.g. Nobilis IB 4/91, D1466, D274, H120, H52 and MA5 (IBV), Nobilis ILT (ILT), Nobilis Marek THV, Rismavac, SB1 and Marexine CA126 (MDV) and Nobilis ND Broiler, Clone 30, Hitchner, LaSota, Nobilis NDC2/Nobilis Newhatch C2 and Nobilis MA5. Such vaccines are

available from Intervet International B.V., Wim de Korverstraat 35, 5831 AN Boxmeer, The Netherlands.

Nobilis Duck Plague is especially suitable for use in combination vaccines for ducks.

Nobilis TRT is especially suitable for use in combination vaccines for turkeys.

5

Preferably, the viral component of the combination vaccine is administered via the same route as the live over-attenuated *Ornithobacterium rhinotracheale* vaccine. *Ornithobacterium rhinotracheale* causes respiratory disease and sometimes severe inflammation reactions in the joints. These inflammation reactions are often seen when *Ornithobacterium rhinotracheale* enters the blood stream. Therefore, a suitable route of administration especially for combating inflammation reactions in the joints is subcutaneous or intramuscular injection of both the bacterial component and a live attenuated poultry virus causing systemic reactions. Examples of viruses, preferred for administration in the combination vaccine according to the invention are live attenuated Avian Reovirus, preferably Nobilis Reo 1133 and 2177, and Marek's virus.

10

15

*Ornithobacterium rhinotracheale* being a mainly respiratory disease, a preferred route however would be oral/nostril/respiratory tract vaccination. A more preferred route of vaccination would be vaccination by spray/aerosol, since in that case, the combined components would most directly reach the respiratory tract and other spots where immunity is most needed.

20

The preferred live attenuated viral components for oral- nostril- or respiratory tract application are those viruses known to cause respiratory disease, specifically Infectious Bronchitis virus, Turkey Rhinotracheitis virus and Newcastle Disease virus.

25

Of these, Nobilis TRT, Nobilis NDC2/Nobilis Newhatch C2 and Nobilis MA5 are the most preferred.

30

Although preferred, both the active components of the combination vaccine do not necessarily have to be administered in a mixed form. The way of administration for each component may depend on the specific properties of each of the components. For example, although administration by injection of such vaccines is also contemplated, the live over-attenuated *Ornithobacterium rhinotracheale* vaccine strains are preferably administered by the inexpensive mass application techniques commonly used for vaccination and well-known in the art, such as spraying/aerosol or drinking water application. It is very well possible that the viral component is preferably given through another route, e.g through parenteral administration.

35

Also other routes of administration are contemplated such as *in ovo* vaccination, of course with the proviso that the specific active component is able to evoke a protective immune response after administration.

- 5 Also, if both the bacterial and the viral component of a certain combination vaccine according to the invention would preferably be administered by spraying, this would not necessarily mean that they are both sprayed through the same nozzle. It could well be possible that, due to differences in viscosity of both the bacterial and viral component, the carriers and/or any excipients, both vaccine components are preferably sprayed through different nozzles.
- 10 Nevertheless, this would of course still lead to the beneficial and synergistic effect that both the viral and the bacterial component of the combination vaccine have: after all, they would both reach the respiratory tract and exert their joined beneficial effect.
- Thus, it goes without saying that such separate administration of both components still gives the surprising effect and therefore is within the scope of the invention.

15

The present invention relies upon the fact that the live attenuated viral component triggers the animal in such a way that the over-attenuated *Ornithobacterium rhinotracheale* vaccine strain stays longer in the respiratory tract and more severely attacks the system. In pathogen-free hosts, the bacterium will disappear within one or two days. Most live attenuated poultry

20

viruses (e.g. vaccine viruses) stay in the respiratory tract for up to a week at least.

Therefore, the advantages of the present invention can also be achieved if the components of the combined vaccine are administered to the birds separated by a small interval of time. The unexpected synergetic effect as described in the present invention will e.g. readily be obtained if the live attenuated viral component is given in the period between seven days before, and

25 two days after the live over-attenuated *Ornithobacterium rhinotracheale* vaccine strain is given.

25

Therefore, combination vaccines of live over-attenuated *Ornithobacterium rhinotracheale* and a live attenuated poultry virus as described in the present invention include those combinations in which the live attenuated viral component is given in the period between

30 seven days before, and two days after the live over-attenuated *Ornithobacterium rhinotracheale* vaccine strain is given, and include those combinations in which the live attenuated viral component and the live over-attenuated *Ornithobacterium rhinotracheale* component are administered at different sites.

30

Merely as an additional example: the advantageous effect of the combination vaccine can also

35 be obtained if one of the components, e.g. the live attenuated poultry virus is administered *in ovo* shortly before hatching, i.e. in the last quarter, preferably at day 18, and the live over-attenuated *Ornithobacterium rhinotracheale* vaccine strain is administered 0-7 days after

35

hatching by e.g. spraying. Preferably, the amount of time between both vaccinations preferably does not exceed 7 days.

5 Preferably, the vaccination with the combi-vaccine is at one day of age. In a more preferred form of this embodiment, the live attenuated poultry virus and the live over-attenuated *Ornithobacterium rhinotracheale* vaccine strain are administered at the same time, if only for ease of application. In an even more preferred form, the live attenuated poultry virus and the live over-attenuated *Ornithobacterium rhinotracheale* vaccine strain are administered by means of spraying.

10 Spraying can be done through a course spray giving visible droplets, or through the use of a nozzle that provides a fine mist. The latter has the advantage that the vaccine more efficiently reaches the lower parts of the respiratory tract.

15 The vaccine according to the invention comprises an effective dosage of the active components, i.e. an amount of immunizing active component that will induce protective immunity in the vaccinated birds against challenge with *Ornithobacterium rhinotracheale* and preferably to the live attenuated viral component of the combination vaccine. (Induction of a protective immunity in the vaccinated birds against the viral component of the combination vaccine is of course desirable, but not in principle necessary).

20 As mentioned above, a protective immune response is an immune response that gives, after challenge, a (statistically significant) decrease in respiratory tract lesion score of equal or more than 50% compared to non-vaccinated animals. In poultry, a higher level of protection against *Ornithobacterium rhinotracheale* would e.g. follow directly from a comparison of lesion scores of vaccinated versus non-vaccinated animals after challenge.

25 The *Ornithobacterium rhinotracheale* component can in principle be administered in a suitable dose of between  $10^3$  and  $10^{10}$  colony forming units. Typically, the *Ornithobacterium rhinotracheale* component will be administered in a dose of 30 between  $10^5$  and  $10^8$  colony forming units.

The live attenuated poultry virus component will usually be given also in a dose of between  $10^3$  TCID<sub>50</sub> and  $10^7$  TCID<sub>50</sub>, although this might of course strongly depend upon the amount of viruses prescribed by the manufacturer of the live attenuated viral component.

The combination vaccine can be used as a primary vaccination, if desired followed by one or more booster vaccinations. The combined vaccine is also suited for incorporation in vaccination programs that also involve the use of other vaccines in a live or inactivated form. Merely as an example, broilers may be vaccinated at one day of age followed by a booster immunization at 10-21 days. Laying stock or reproduction stock may be vaccinated at 1-10 days followed by booster vaccinations at 26-38 days and 16-20 weeks.

In a more preferred form, the combination vaccine comprises a combination of an over-attenuated *Ornithobacterium rhinotracheale* and Newcastle Disease virus, even more preferably NCD Clone 30, obtainable from Intervet International B.V., Wim de Korverstraat 35, 5831 AN Boxmeer, The Netherlands.

In an still even more preferred form, the combination vaccine comprises a combination of an over-attenuated *Ornithobacterium rhinotracheale* and Newcastle Disease virus NDC2, as deposited at the CNCM of the Institut Pasteur, 25 Rue du Docteur Roux, Paris, France under Accession number I-1614.

In another more preferred form, the combination vaccine comprises a combination of an over-attenuated *Ornithobacterium rhinotracheale* and a live attenuated Turkey Rhinotracheitis virus, even more preferably Nobilis TRT, obtainable from Intervet International B.V., Wim de Korverstraat 35, 5831 AN Boxmeer, The Netherlands.

In again another more preferred form, the combination vaccine comprises a combination of an over-attenuated *Ornithobacterium rhinotracheale* and a live attenuated Infectious Bronchitis virus, even more preferably Nobilis MA5, obtainable from Intervet International B.V., Wim de Korverstraat 35, 5831 AN Boxmeer, The Netherlands.

In still another more preferred form, the combination vaccine comprises a combination of an over-attenuated *Ornithobacterium rhinotracheale* and a live attenuated Avian Reovirus, even more preferably Nobilis Reo 1133 or 2177, obtainable from Intervet International B.V., Wim de Korverstraat 35, 5831 AN Boxmeer, The Netherlands.

The invention also relates to combination vaccines comprising, in addition to the two active components defined above, one or more additional antigens derived from a virus or micro-organism pathogenic to poultry or genetic information encoding said antigen.

More preferably, the virus or micro-organism is selected from the group consisting of Infectious Bronchitis virus, Infectious Bursal Disease (Gumboro), Chicken Anaemia agent,

Avian Reovirus, *Mycoplasma gallisepticum*, Turkey Rhinotracheitis virus, *Haemophilus paragallinarum* (Coryza), Chicken Poxvirus, Avian Encephalomyelitisvirus, Duck Plague virus, Egg Drop syndrome virus, Infectious Laryngotracheitis virus, Herpes Virus of Turkeys, *Eimeria* species, *Ornithobacterium rhinotracheale*, *Pasteurella multocida*, *Mycoplasma synoviae*, *Salmonella* species and *E. coli*.

The vaccine according to the invention can be prepared and marketed in the form of a suspension, or in a lyophilized form and additionally contains a pharmaceutically acceptable carrier customarily used for such active components. Carriers include stabilisers, diluents, preservatives and buffers.

Suitable stabilisers are for example SPGA, carbohydrates (such as dried milk, serum albumin or casein) or degradation products thereof. Suitable buffers are for example alkali metal phosphates. Suitable preservatives are thimerosal, merthiolate and gentamicin. Diluents include water, aqueous buffers (such as buffered saline), alcohols and polyols (such as glycerol).

If desired, the vaccine according to the invention may contain an adjuvant. Suitable compounds or compositions for this purpose include alum hydroxide, -phosphate, or -oxide, oil-in-water or water-in-oil emulsions based upon for example mineral oil, such as Bayol F® or Marcol 52® or a vegetable oil such as vitamin E acetate, and saponins.

Preferably, both the bacterial and the viral component of the combination vaccine according to the invention are packaged in a mixed form in the same container. They could be freeze-dried or stored at low temperature thereafter.

The present invention also contemplates a kit format which comprises a packaged multi-container unit having containers each comprising one of the active components as defined above. The kit may additionally comprise a container with a carrier for one or both of the active components, in which case the carrier preferably comprises an adjuvant. Such a kit is e.g. advantageous in case the active components are preferably not freeze-dried or premixed together, or when the active components are preferably administered separated in place and/or time.

The active components of the kit according to the invention, i.e. the live over-attenuated *Ornithobacterium rhinotracheale* strain and the live attenuated poultry virus, and possibly a third and further active components, can be mixed prior to vaccination, or administered

separately, at different administration sites and/or different moments of administration with the proviso's indicated above.

- 5 Still another embodiment of the present invention relates to the use of a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus for the manufacturing of a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*.

- 10 Also, another embodiment relates to the use of a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus for the manufacturing of a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*, wherein the live over-attenuated *Ornithobacterium rhinotracheale* strain and the live attenuated poultry virus are administered simultaneously, separately or sequentially. Simultaneous administration is administration of the live over-attenuated *Ornithobacterium*  
 15 *rhinotracheale* strain and the live attenuated poultry virus at the same moment in time, preferably injected as a mixture, sprayed as a mixture sprayed from one nozzle or given as a mixture in the drinking water. Separate administration is administration of the live over-attenuated *Ornithobacterium rhinotracheale* strain and the live attenuated poultry virus from  
 20 two different injection sites or, when sprayed, e.g. from two different nozzles, preferably at the same moment in time. Sequential administration is administration during which the live over-attenuated *Ornithobacterium rhinotracheale* strain and the live attenuated poultry virus are administered at different moments in time. The conditions for the various moments of administration are discussed above.

- 25 Again another embodiment of the present invention relates to methods for the preparation of a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*. Such methods comprise the admixing of a live over-attenuated *Ornithobacterium rhinotracheale* strain, a live attenuated poultry virus and a pharmaceutically acceptable carrier.  
 30 Such methods may also and/or alternatively comprise the packaging of a live over-attenuated *Ornithobacterium rhinotracheale* and a live attenuated poultry virus in separate containers that form part of a kit.



## **EXAMPLES**

### **Example 1**

#### **Preparation of live over-attenuated *Ornithobacterium rhinotracheale* strains.**

- 5 Two commonly known attenuation targets, viz. *purD* and *recA*, were selected to create mutants.

Since serotype A strains are the most predominant and is also known to provide cross-protection against infections with other serotypes, this serotype was used to create the mutants.

10

#### **Bacterial strains and plasmids**

*Escherichia coli* host strains TOP10 and TOP10F' were purchased from Invitrogen (Carlsbad, CA). pUC19 was derived from Clontech laboratories (Palo Alto, CA).

- 15 An *Ornithobacterium rhinotracheale* strain serotype A (*OR-7* (O95264 95.2932) Van Empel, P.C.M.; Molecular identification of *Ornithobacterium rhinotracheale*; (1998) ISBN 90-393-1574-4. Page 45.) was used as host strain for electroporation and homologous recombination. Chromosomal DNA of *Ornithobacterium rhinotracheale* strain 3263/91 (see above) was used as PCR template for making deletion constructs.

#### **20 Culture media, buffers and antibiotics**

Luria-Bertani (LB) broth and terrific broth (TB, Sambrook, J., E.F. Fritsch, and T. Maniatis. Molecular cloning: a laboratory manual. (1989) Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.) were used for liquid growth of bacterial *Escherichia coli* strains. For *Ornithobacterium rhinotracheale* (*OR*) Todd Hewitt (Becton Dickinson, Franklin Lakes, NJ, non irradiated) was used. For *OR* on plates, blood agar or Todd Hewitt agar supplemented with 4% sheep blood was used. LB agar plates were poured from melted LB-agar, which was cooled to 45°C and then the appropriate antibiotics were added.

25

#### **Oligonucleotides**

- 30 All oligonucleotides were obtained from Life Technologies™ Gibco BRL (Paisley, UK) and are indicated in Table 1. Their location in the sequence is shown in figure 2A and B.

#### **PCR amplification**

- 35 PCR amplification was performed where the PCR mixture for cloning the *recA* and *purD* genes consisted of 40 U/ml of SuperTaq plus; 1x SuperTaq PCR buffer; 80 µM (each) of dATP, dCTP, dGTP and dTTP (HT BioTechnology Ltd., Cambridge, UK); 0.4 µM of both oligonucleotides and 1 µl of chromosomal DNA as template. Samples were analyzed on an

agarose gel. For the overlap extension PCR (OE-PCR) the first PCR products were diluted a hundred fold and mixed 1:1. Then PCR was performed with the outer primers as described below.

The colony PCR method was performed as follows: part of the colony was picked from an agar plate and transferred into a PCR tube. The PCR mixture was added, containing: 10 U/ml of SuperTaq DNA polymerase; 1x SuperTaq PCR buffer; 80  $\mu$ M (each) of dATP, dCTP, dGTP and dTTP (HT BioTechnology Ltd., Cambridge, UK) and 0.4  $\mu$ M of both oligonucleotides. Samples were analyzed on an agarose gel.

10 Table 1. Oligonucleotides (primers) used for preparation of insertion-deletion constructs.

Primer Name	Sequence (5' $\rightarrow$ 3')	Features
<i>PurD</i> -F13	CTTAAGCTTGGATCCTTGTGGCGTGGCTTTAG	<i>HindIII</i> site (underlined)
<i>PurD</i> -OE-R	CTTCTAGCGTAGCGCCAGATCTCATTGTTCGGT TCCAGCGTTTCC	<i>BglIII</i> site (underlined) and overlap with <i>PurD</i> -OE-F (italics)
<i>PurD</i> -OE-F	GAGATCTGGCGCTACGCTAGAAGAAGCC	<i>BglIII</i> site (underlined) and overlap with <i>PurD</i> -OE-R (italics)
<i>PurD</i> -R8	CTTAAGCTTCAGTGGAGCGGCAGATACAGAG	<i>HindIII</i> site (underlined)
<i>RecA</i> -F6	CTTAAGCTTGGAGCGTGTAGTGCTCGCCATCG	<i>HindIII</i> site (underlined)
<i>recA</i> -OE-R	ACCGCACGCACGAGATCTCGGGCTTTGTGCGCCC ATCATCATCAC	<i>BglIII</i> site (underlined) and overlap with <i>recA</i> -OE-F (italics)
<i>recA</i> -OE-F	CGAGATCTCGTGCGTGCGGTATTGAAAG	<i>BglIII</i> site (underlined) and overlap with <i>recA</i> -OE-R (italics)
<i>RecA</i> -R5	CTTAAGCTTCCCAGCCAATTCGGCTCGTTTCAC	<i>HindIII</i> site (underlined)

#### Nucleotide sequence analysis

The nucleotide sequences of the *purD* and *recA* genes and their surroundings were determined.

15 Sequencing was performed using a cycle sequencing protocol on a Perkin Elmer 9700 DNA thermal cycler with big dye terminators: approx. 50 ng of PCR product, 2  $\mu$ l terminator ready reaction mix (Perkin Elmer, CA) 2.4 pmoles of primer, 6  $\mu$ l of buffer (200 mM Tris-HCl pH8.5; 5mM MgCl<sub>2</sub>) and aqua dest in a total of 20  $\mu$ l.

#### 20 DNA manipulations

For sequencing, the *purD* and *recA* genes and their flanking regions were amplified by PCR and cloned into a general purpose cloning vector.

### Electroporation of *Ornithobacterium rhinotracheale*.

*Ornithobacterium rhinotracheale* was made electro-competent by growing in Todd-Hewitt broth until an OD 600 nm of 0.3. During further manipulations, the cells were kept on ice.

- 5 The culture was washed twice with ice-cold water and once with ice cold 10 % glycerol. This suspension was centrifuged again and the cells were resuspended in 0.5 ml of 10% glycerol and stored on ice until use.

- Electroporation was performed using a BTX ECM630 Genepulser (San Diego, CA). 1–2 µg of plasmid DNA was mixed with 50 µl of electro competent *Ornithobacterium rhinotracheale* cells in a 2 mm cuvette. A pulse of 20–24 msec. was delivered using 2500V; HV capacity; 25 µF and 800 Ω. After electroporation, 1 ml of TH broth was added and the cells were recovered for 2 hours at 37°C and 100 RPM. Viability was checked on blood agar plates. Antibiotic resistant colonies were selected on TH agar with 4 % blood and antibiotic at a selective concentration.

15

### UV-sensitivity of *recA* recombinants.

- In order to check a recombination in the *recA* gene, its sensitivity to UV was tested. Bacteria were plated onto blood agar and UV irradiated with a wavelength of 365 nm at a distance of about 10 cm. Different exposure times were tested, ranging from 0 to 120 seconds. Plates were then incubated and growth of *recA*-mutant colonies were compared to growth of *purD*-mutant colonies and colonies of the wild type strain.

20

## RESULTS

### Sequencing *recA* and *purD*.

- 25 To obtain the sequence of the *purD* and *recA* genes of *OR*, degenerate primers were developed based on conserved regions of the genes derived from closely related bacteria. By means of genome walking, the flanking regions were determined. The complete sequences and the relevant features are shown in figure 2A for *purD* and figure 2B for *recA*.

### 30 Construction of the *recA* and *purD* deletion and insertion-deletion constructs.

- The strategy for construction an insertion-deletion mutant is depicted in Figure 1. For the *purD* mutant the procedure was as follows. First, PCR on genomic *Ornithobacterium rhinotracheale* DNA was performed using 2 primer sets *purD*-F13 (+*HindIII* site) with *purD*-OE-R (upstream *purD*) and *purD*-OE-F with *purD*-R8 (+*HindIII* site) (downstream *purD*) to amplify the flanking regions of the *purD* gene. Overlap-extension PCR was performed with the *purD*-F13 and the *purD*-R8 primers using both *purD* PCR fragments as templates since the OE-R and OE-F primers overlap. This results in PCR fragments that contain both flanking

35

regions of the *purD* gene and therefore the biggest part of the *purD* gene is deleted. These fragments were digested with *HindIII* and cloned into pUC19. Then an antibiotic resistance marker present on a 1.2 kb *BamHI* fragment was introduced in the *BglIII* site that was included in the overlap sequence of the OE-F and OE-R primers.

- 5 A similar approach was followed for construction of the *recA* insertion-deletion construct. Primers *recA*-F6 and *recA*-OE-R (upstream *recA*), and *recA*-OE-F and *recA*-R5 (downstream *recA*) were used in PCR for the flanking regions of the *recA* gene. OE-R and OE-F primers overlap and have an additional *BglIII* site for insertion of the antibiotic resistance marker. The constructs were verified, plasmid DNA was isolated and used to electroporate
- 10 *Ornithobacterium rhinotracheale*.

#### Verification of the mutation in recombinants.

After electroporation, colonies were obtained for both the *recA* and *purD* constructs, that showed antibiotic-resistance.

- 15 Mutants for both genes were confirmed by PCR. The *recA* mutant clones were additionally confirmed by testing for their sensitivity to UV irradiation; the *purD* mutants and the parent strain were able to survive UV exposure times for more than 40 sec whereas the *recA* mutant died after 10 sec of irradiation.

20

#### Example 2

**Comparison of induction of protection of over-attenuated *Ornithobacterium rhinotracheale* strains.**

Vacc. 2 weeks 2*10 <sup>7</sup> CFU/ml	Prim. Day 23	Chall. Day 28 3*10 <sup>8</sup> CFU	Resp. score.
<i>O.r.</i> RecA	NDV	100 ml spray	46
<i>O.r.</i> PurD	NDV	100 ml spray	35
--	NDV	100 ml spray	51

25

Table 2: comparison of level of protection of over-attenuated mutants. Vaccine strains were applied by spraying. Each group consisted of 15 chickens. Challenge was done with wild-type. Respiratory scores are given as a percentage of the maximally possible score of the challenged group.

30

The affected organs of the birds were scored in a numerical scoring system:

Thoracic air sacs (TAS): 0 = no abnormalities, 1 = one air sac seriously affected by fibrinous airsacculitis or limited pinhead-sized foci of fibrinous exudate in both air sacs, 2 = both air sacs seriously affected by fibrinous airsacculitis.

5 Abdominal air sacs (AAS): 0 = no abnormalities, 1 = pinhead-sized foci of fibrinous exudate or slight diffuse airsacculitis, 2 = severe fibrinous airsacculitis.

Pneumonia: 0 = no abnormalities, 1 = unilateral pneumonia, 2 = bilateral pneumonia.

Trachea: 0 = no abnormalities, 1 = some exudate in the tracheal lumen, 2 = lumen of trachea filled with exudate.

10 Table 2 shows the result of a vaccination experiment in which two weeks old chickens were vaccinated by spray-vaccination of  $2 \times 10^7$  CFU in 100 ml spray solution of a RecA-mutant and a PurD-mutant of *Ornithobacterium rhinotracheale*. Priming was done with an NDV Lasota vaccine strain at day 23, and challenge with a wild-type *Ornithobacterium rhinotracheale* strain was done at day 28. The level of protection was determined by counting the trachea and  
15 lung lesions.

The figures of the respiratory lesion score (see column resp. score in table 2) are to be interpreted as follows: the maximal possible lesion score is taken as 100% score. Non-vaccinated animals show a lesion score of 51%, and "vaccinated" animals have a lesion score  
20 of 46% (RecA) or 35% (PurD). This means that "vaccination" with RecA or PurD gives a level of protection of 8% ( $100\% - (46 \times 100\% / 51)$ ) or 30% respectively.

A protective immune response, as defined above, is an immune response that gives, after challenge, a (statistically significant) decrease in respiratory tract lesion score of equal or more than 50% compared to non-vaccinated animals.

25 It follows from table 2 that the respiratory lesion score of the groups that received vaccination with a mutant of *Ornithobacterium rhinotracheale* does not significantly differ from the respiratory lesion score found with the non-vaccinated animals.

Conclusion: a RecA-mutant and a PurD-mutant of *Ornithobacterium rhinotracheale*, if given as such, are not capable to raise a protective immune response in chickens.

**Example 3**

**Comparison of induction of protection of over-attenuated *Ornithobacterium rhinotracheale* strains after priming and boosting.**

Vacc. Day 1 $2 \times 10^7$ CFU/ml strain RecA	Boost 2 weeks $2 \times 10^7$ CFU/ml strain-RecA	Prim. Day 23	Chall. Day 28 A: $4 \times 10^7$ Spray.	Resp. score.
--	--	NDV	A	27
Nostr./conj.0.2 ml	Spray 100 ml	NDV	A	19 <sup>b</sup>

5

**Table 3:** comparison of level of protection obtained with over-attenuated mutants after a first administration followed by a booster administration. Each group consisted of 12 chickens. Respiratory scores are given as a percentage of the maximally possible score of the challenged group. Challenge was done with wild-type

10

(b) not statistically different from non-vaccinated control.

15

Table 3 shows the result of a vaccination experiment in which chickens have been vaccinated first in the nostrils and then boosted after two weeks by spray-vaccination using a RecA-mutant of *Ornithobacterium rhinotracheale*. Priming was done with an NDV Lasota vaccine strain at day 23, and challenge with a wild-type *Ornithobacterium rhinotracheale* strain was done at day 28. The level of protection was determined by counting the respiratory lesions.

20

As can be seen in the last column, the respiratory lesion score of the groups that received two subsequent vaccinations with a RecA-mutant of *Ornithobacterium rhinotracheale* does not significantly differ from the respiratory lesion score found with the non-vaccinated animals. Conclusion: a RecA-mutant of *Ornithobacterium rhinotracheale* is not capable, even when given twice, to raise a protective immune response in chickens.

**Example 4****Induction of lesions by over-attenuated mutants and wild-type strains.**

Prim. Day 23	Chall. Day 28 100 ml spray	Resp. score.
NDV	RecA $7 \times 10^7$	22
NDV	PurD $2 \times 10^8$	24
NDV	Wild-type	59

- 5 **Table 4:** comparison of induction of lesions by over-attenuated mutants and wild-type strains. Each group consisted of 12 chickens. Respiratory scores are given as a percentage of the maximally possible score of the challenged group.

10 Table 4 shows the results of challenge with both an RecA-mutant, a PurD-mutant and a wild-type strain of *Ornithobacterium rhinotracheale*. No vaccination took place prior to priming and challenge. As can be seen from the last column, the mutants give a respiratory lesion score that is less than half the respiratory lesion score found after challenge with the wild-type strain. This clearly shows that the mutants behave mild, even without prior vaccination, and even after priming with a relatively "hot" NDV-strain, i.e. the Lasota strain.

15

**Example 5****Comparison of protection of *Ornithobacterium rhinotracheale*/NDV combination vaccines after subsequent challenge.****Experimental design**

- 20 A 125 1-day-old SPF broiler chickens were used. At 1-day-old all chickens were spray vaccinated with live attenuated NDV and subsequently groups of 25 chickens were aerosol vaccinated with either of the mutant strains *recA* and *purD* or the wild type strain 3263/91 (serotype A) or were left unvaccinated (challenge controls). At 9 days after vaccination 10
- 25 chickens of each group were sacrificed for post-mortem examination to assess the safety of each vaccine. The remaining chickens (15 per group) were spray treated with ND-Lasota at 25-days of age and aerosol challenged with wild-type *Ornithobacterium rhinotracheale* 3263/91 (serotype A) at 31 days of age. Seven birds of each group were additionally challenged intravenously. At 38 days of age the chickens were killed and post-mortem examined to assess the efficacy of each vaccine.

In addition to a non-*Ornithobacterium rhinotracheale*-vaccinated challenge control group, a non-*Ornithobacterium rhinotracheale*-vaccinated and non-challenged control group was used as NDV-control group.

The detailed experimental design and treatment schedule is shown in Table 5.

5

#### *Ornithobacterium rhinotracheale* vaccine cultures

For aerosol vaccination with the wild type *Ornithobacterium rhinotracheale* 3263/91 (serotype A), *recA* and *purD*, fresh cultures in Todd-Hewitt broth containing  $6.7 \times 10^8$  CFU/ml,  $3.9 \times 10^7$  CFU/ml and  $2.6 \times 10^8$  CFU/ml, respectively, were used. The dose for aerosol

10

#### ND suspensions

NDV-strain NDC2 has been deposited at, and is obtainable as Nobilis NDC2/Nobilis Newhatch C2, or from the CNCM of the Institut Pasteur, 25 Rue du Docteur Roux, Paris, France under Accession number I-1614. ND-Lasota was obtained from Intervet Int. B.V. (see above). The dose for NDC2 was  $10^{6.4}$  EID<sub>50</sub> per animal (corresponding to 2 ml per bird) and for ND-Lasota  $10^{6.6}$  EID<sub>50</sub> per animal (corresponding to 3 ml per bird), using a 1 L spray can.

15

#### *Ornithobacterium rhinotracheale* challenge culture

For aerosol challenge a fresh culture of wild-type *Ornithobacterium rhinotracheale* strain 3263/91 (serotype A) in Todd-Hewitt broth containing  $4.3 \times 10^8$  CFU per ml (as determined by plate counting) was used.

20

Animals: 125 1-day-old SPF Hybro broilers were used.

25

#### Grouping and dosing (vaccination / challenge)

Chickens were randomly divided (as they came to hand) over the different isolators and were treated as shown in Table 5.

NDC2 (1-day-old) or ND-Lasota (25-day-old) were given by spray using a 1 L spray can: 2 ml per bird or 3 ml per bird, respectively. The birds remained in the spray for 10 minutes with the air-circulation closed. At 1-day-old the birds were first treated with NDC2 and subsequently with *Ornithobacterium rhinotracheale*.

30

*Ornithobacterium rhinotracheale* vaccine (1-day-old) as well as *Ornithobacterium rhinotracheale* challenge (31-day-old) was given by aerosol at a dose of 100 ml per isolator using a paint sprayer.

35

After the aerosol challenge 7 birds of each group were additionally challenged intravenously at a dose of 1 ml into the wing vein.



Post mortem investigations

Birds that died during the experiment were subjected to a post-mortem examination to determine the cause of death.

- 5 At 10-day-old (9 days after vaccination) 10 birds per group were sacrificed and subjected to a post-mortem examination to assess the safety (attenuation) of the (vaccine) strains compared to the wild type. At 38 days of age (7 days after challenge) the (remaining) challenged birds were killed and also subjected to a post-mortem examination to assess the efficacy of the different strains.
- 10 The affection rates of the groups are given as a percentage of the maximal possible lesion score of the group.

Statistical analysis

- 15 The total lesion scores per group (compared to the controls) were analysed two-sided using the non-parametric Mann-Whitney U test. The level of significance was set at 0.05.

Results and conclusion

- 20 Concurrent aerosol vaccination at 1-day-old with live attenuated NDV and wild type *Ornithobacterium rhinotracheale* induced unacceptable lesions after vaccination but also (as a result of that) induced a strong protective response as evidenced after challenge where this group had the least post-mortem scores. Apparently, the initial lesions caused by the vaccination had already disappeared at time of challenge.

- 25 Concurrent aerosol vaccination at 1-day-old with live attenuated NDV and *recA* or *purD* mutant *Ornithobacterium rhinotracheale* strains demonstrated a good level of safety as well as efficacy.

- 30 As follows from table 5, concurrent aerosol vaccination of 1-day-old SPF broilers with live attenuated NDV and *recA* or *purD* mutant strains appeared safe and to induce a good level of immunity against challenge with wild type *Ornithobacterium rhinotracheale*.

Example 6

**Comparison of protection of *Ornithobacterium rhinotracheale*/MA5 combination vaccines after subsequent challenge.**

In this experiment, Infectious Bronchitisvirus type MA5 was used as the live attenuated viral component. Additionally, the experiment with *Ornithobacterium rhinotracheale*/NDV combination vaccines was repeated.

The experimental set-up of this Example was largely identical to that of Example 5. Where

5 different numbers of animals or different moments of vaccination or challenge apply, this is indicated in table 6. This table also gives an overview of the vaccines used and the

---

vaccination schedule, as well as the level of protection obtained.

MA5 suspensions: live attenuated Infectious Bronchitisvirus (IBV) type MA5 (Intervet International B.V., Wim de Korverstraat 35, 5831 AN Boxmeer, The Netherlands) was used

10 at a concentration of  $5.5 \log^{10}$  EID<sub>50</sub> per animal, and applied by spraying.

### Conclusion

As follows from table 6, concurrent aerosol vaccination of 1-day-old SPF broilers with IBV MA5 and PurD-mutant strains induces a good level of immunity against challenge with wild-  
15 type *Ornithobacterium rhinotracheale*. Additionally it follows from table 6, that concurrent aerosol vaccination of 1-day-old SPF broilers with NCD and PurD-mutant strains induces a good level of immunity against challenge with wild-type *Ornithobacterium rhinotracheale* as was also demonstrated in Example 5.

table 5

group	no. of chickens	Treatment			Results	
		vaccination at day 1	challenge at day 25	challenge at day 31	% of max airsac score at day 10 (safety)	% of max airsac score at day 38 (efficacy)
1	25	NDV	RecA aerosol	NDV	WT-OR aerosol	2.5 <sup>b</sup>
2	25	NDV	PutD aerosol	NDV	WT-OR aerosol	2.5 <sup>b</sup>
3	25	NDV	WT-OR aerosol	NDV	WT-OR aerosol	10 <sup>b</sup>
4	25	NDV		NDV	WT-OR aerosol	47
5	25	NDV		NDV	NDV	2

<sup>b</sup> Significantly different ( $p < 0.05$ ) compared to the controls (group 1) using two-sided Mann-Whitney U test

table 6

group	no. of chickens	Treatment			Results
		vaccination at day 1		challenge day 35	
1	15		PurD aerosol	NDV	WT-OR aerosol
2	15	NDV	PurD aerosol	NDV	WT-OR aerosol
3	15	NDV		NDV	WT-OR aerosol
4	15	MA5		NDV	WT-OR aerosol
5	15	MA5	PurD aerosol	NDV	VT-OR aerosol
					no reduction
					54% <sup>b</sup>
					no reduction
					no reduction
					50% <sup>b</sup>

<sup>b</sup> Significantly different ( $p < 0.05$ ) compared to the controls (group 11) using two-sided Mann-Whitney U test

### Legend to the figures

Figure 1: cloning strategy for making mutation constructs of *recA* and *purD* genes in OR. Overlap extension PCR was used to create a deletion in the open reading frame of the gene of interest and these fragment were cloned in pUC19. Subsequently an antibiotic resistance marker gene (BamHI fragment) was cloned into the BglII site of the gene of interest (insertion-deletion constructs).

---

Figure 2A: Sequence of *purD* and localization of primers used for cloning and verification of mutants.

Figure 2B. Sequence of *recA* and localization of primers used for cloning and verification of mutants

## Claims

- 1) A combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*, characterized in that said combination vaccine comprises a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus.
- 2) A combination vaccine according to claim 1, characterized in that said live attenuated poultry virus is Infectious Bronchitis virus, Newcastle Disease virus, Turkey Rhinotracheitis virus, Marek's virus or Avian Reovirus.
- 3) A combination vaccine according to claim 1, characterized in that said live attenuated poultry virus is Infectious Bronchitis virus, Newcastle Disease virus or Turkey Rhinotracheitis virus.
- 4) A combination vaccine according to claim 1, characterized in that said live attenuated poultry virus is Newcastle Disease virus.
- 5) A combination vaccine according to claim 1, characterized in that said live attenuated poultry virus is Newcastle Disease virus type NDC2.
- 6) A combination vaccine according to claim 1, characterized in that said live attenuated poultry virus is Turkey Rhinotracheitis virus.
- 7) A combination vaccine according to claim 1, characterized in that said live attenuated poultry virus is Infectious Bronchitis virus.
- 8) A combination vaccine according to claims 1-7, characterized in that said live over-attenuated *Ornithobacterium rhinotracheale* has a mutation, said mutation preferably being a deletion, in the *purD*-gene or the *recA*-gene.
- 9) A combination vaccine according to claims 1-8, characterized in that said combination vaccine comprises an additional antigen derived from a virus or micro-organism pathogenic to poultry or genetic information encoding said antigen.
- 10) A combination vaccine according to claim 9, characterized in that the virus or micro-organism is selected from the group consisting of Infectious Bronchitis virus,

Infectious Bursal Disease (Gumboro), Chicken Anaemia agent, Avian Reovirus, *Mycoplasma gallisepticum*, Turkey Rhinotracheitis virus, *Haemophilus paragallinarum* (Coryza), Chicken Poxvirus, Avian Encephalomyelitisvirus, Duck Plague virus, Egg Drop syndrome virus, Infectious Laryngotracheitis virus, Herpes Virus of Turkeys, *Eimeria* species, *Ornithobacterium rhinotracheale*, *Pasteurella multocida*, *Mycoplasma synoviae*, *Salmonella* species and *E. coli*.

- 11) Use of a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus for the manufacturing of a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*.
- 12) Use of a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus for the manufacturing of a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*, wherein the live over-attenuated *Ornithobacterium rhinotracheale* strain and the live attenuated poultry virus are administered simultaneously, separately or sequentially.
- 13) Method for the preparation of a combination vaccine according to claims 1-10, characterized in that said method comprises the admixing of a live over-attenuated *Ornithobacterium rhinotracheale* strain, a live attenuated poultry virus and a pharmaceutically acceptable carrier.
- 14) A vaccination kit for the immunization of poultry against *Ornithobacterium rhinotracheale*, characterized in that said kit comprises
  - a) a live over-attenuated *Ornithobacterium rhinotracheale* strain and
  - b) a live attenuated poultry virus and
  - c) optionally a pharmaceutically acceptable carrier for the component under a and/or b.
- 15) A vaccination kit according to claim 14, characterized in that the carrier comprises an adjuvant.

## Abstract

The present invention relates to a combination vaccine for the protection of poultry against *Ornithobacterium rhinotracheale*, to the use of a live over-attenuated *Ornithobacterium rhinotracheale* strain and a live attenuated poultry virus for the manufacturing of such a combination vaccine, to methods for the preparation of said combination vaccine and to vaccination kits for the immunization of poultry against *Ornithobacterium rhinotracheale*.

---





Figure 1. Overlap-extension-PCR fragment with purD deletion

Overlap-extension-PCR fragment with recA deletion




Figure 2A.

1 GTTCGACCAA ACGGCTTGT GTGCGGTGAA ACATAGCACT CCTTGTGGCG TGGCTTTAGA TGATGATATT TTGCAAGCGT  
>>.....F5.....>> CTTAAGCTTGGA>>.....F13.....>>

-----  
HindIII

81 ACCAAAAAGC ACACGACTGC GACCCGATTT CGATTTTGG TGGCATTGTA ACTTTTAATA AAAAAGTAAC AAAAGCAGTG  
161 GCAGAAAAAT GTAACGAGAT TTTCTTGAA ATCGTTGCTG CACCGAGCTT TGAGCCAGAG GCTTTGGAAG TTTTGTCTAA  
241 AAAGAAAAAT TTGCGCGTGA TTGAAGTTAA AAATCCATTA AGCGATAAAA TGCAACTCGT GCAAGTAGAT GCGCGATTGC  
321 TCGTGCAAGA AATCGACAAA TCGTTTAGCA ATGATTTTAA AGTAGTAACC GAAAAACAAC CTACCGAAAA GCAACTTTCT  
401 GATTTGGAAT TTGCCATGAA AGTAGTGAAG CATGTAAAGA GCAATGCCAT CGTGGTTGCC ACAAACGGAC AAGCTCTAGG  
481 CGTGGGCACA GCGGAGACTA ATCGTATTTG GGCAGCACAG CAGGCGATTG AGCGTGCAAA GGAAAAACA CAAGAAAATC  
561 TAGTTTGGC TTCCGATGCC TTTTCCCAT TCAGAGATGT GGTAGATTAT GCAGCACAAG AAGGCATTAC AGCCTTGATT  
641 CACCCAGGAG GAAGCATGCG CGACCAAGAG AGCATAGACG CGGCTAATGA ACACGGAATC CCGATGATCA TCAGCGGTAT  
721 GAGACATTTC TTACATTAA TCAAAAAATC TAAACAATAA TTATCAATAA TTCTAAAACA CAATAAGTAT GAATGCAAT  
>>...purD...>

801 GATTACAAAA AAATACTCAT CGTAGGAAAC GCGCAAGAG AACACGCCAT CGGGTGGAAT ATTAACAAG ACCACCCTTC  
>.....purD.....>

881 TTGCGAGCTT TTCTTTGCGC CAGGAAACGC TGAACCGAA CAAATTGGAA AAAACATCGT AGCTGAATCT AATTATGGCT  
>.....purD.....>  
<<.....OE-R.....<<AGATCTGGCGCTACGCTAGAAG

-----  
BglIII

961 TAATGCTTTT TGCTCAACAA AATGATATAG ACTTAACGAT TGTAGGTCCA GAAGCAGAAT TGGTAGAAGG TATTGTAGAC  
>.....purD.....>  
1041 TTGTTTGAAT CCAATCAATT AAGAATTTT GGTCCAGATA AGCGTGCGGC TAAATTGGAA GGCAGCAAGG CTTTGTCCAA  
>.....purD.....>  
1121 AGATTTTATG GAGAAATACG GCGTGCGCAC GGCTTTTGCC AAAAGTTTCA ACAATTTTGT AGACGCTAGA GATTATGTAA  
>.....purD.....>  
1201 AAGAGCTCAC GCAATTCCTT ATCGTGATCA AAGCCAGTGG CTTGGCAGCA GGAAAAGGTG TGATCATCGT GCACNTACAA  
>.....purD.....>  
1281 CTTGAAGCCG AAATCTACTT GCGCAAAATC ATGGAAGACA AAACCTTTGG CGAAGCAGGC AACGAGGTCG TAATCGAGGA  
>.....purD.....>  
1361 ATACTTAAAA GGTGTGGAAG TTTCTGTGCT TTCTATCTTT AACCATAAAG AAATTAAAAC TTTCTTGCCT GTAAAAGACC  
>.....purD.....>  
1441 ACAAGAAAAAT CGGAAAAGGC GAAACAGGAC TCAACACGGG CGGAATGGGC GTAGTGGCTC CTAACCCGCA TTTTACCGAT  
>.....purD.....>  
1521 GAGCACATGA AGGAGTTTGA GAAAAACATT TTGCTCCCAA CACAAAAGG GCTCTTGGCA GAAAAATGC ATTTTGCAGG  
>.....purD.....>  
1601 CATTATTTTC TTTGGGCTTA TGATTACCGA GCATGGTATT TATCTATTGG AATACAACAT GCGATTGGC GACCCAGAAA  
>.....purD.....>  
1681 CCGAAGCACT TTGCTTTT ATGGAGAATG ATTTAGTAGC CCTCATCGAT TCCGCAATAC ACCAGCAAGA CATTGAACCT  
>.....purD.....>  
1761 AAATGGAAAA ACGAACATGC TTGCTGTGTA GTAATGGCGA GCGGTGGCTA CCCAGGCACT TACGAAACTG GTTTTGAAAT  
>.....purD.....>

1841 CCGAGGATTG AACAAAGTTG ATGTTCCCGT ATTTATTGCA GGAGCCAGAG AAGAAAGTGG AAAAATCTAC ACCACAGGCG  
>.....purD.....>

1921 GGCGCGTGCT CAATGTGGTG GGAAGTGGCG CTACGCTAGA AGAAGCCAGA AAAGTGGCTT ACGAAAATAT CCATAAAATC  
>.....purD.....>

GAGATCTGG>.....OE-F.....>

-----  
BglIII

2001 AATTTTGATT ATGAATATTA TCGCGAAGAC ATCGGGAAGA TATAATCTCG CTGATTTTAA ACCAAAACAT ATTTAAAAAC  
>.....purD.....>

2081 GCTTTTGTTA CTTTTATAAA CAAAGGCGTT TTTCTATTTT TGTGCCACTA TAACATGATT TAACCCATGA AAAAAATACT

2161 AAAAAATCTC ATTTTCTTAC TGCTCATTC CTGGGTTTAT GCCCTGATTT TAATCTTTAT AAATCCACCT ATCACCATTA

2241 CACAGCTGAG CAATTTATCT TATGGTTTCT CCAGAACACA GCTCGCTTAT GATGAAATTC CGGCTAGTGC TAAATGGGCT

2321 GTAATTGCAG CAGAAGACCA GAATTTTGCC ATTCATAATG GCTTTGATTT TAAAGAAATT AAAACCGCCT ACGAGAAAAA

2401 CAAAGCGGGC AAGAAATTGC GTGGCGGGAG CACCTTTTCG CAACAACTG CCAAAAATGT ATTTTGTGG CAAGGGCGCA

2481 CTTGGATTAG AAAAGGATTG GAAACCTACT GCACCTTTAT CATCGAAACG CTGTGGAGCA AGGAGCGTAT TTTGCAAGTT

2561 TACCTCAACA ATGCCGAAAT GGGCAAAGGC GTTTATGGCA TAGAGGCAGC GGCGCAATAT TATTTTAAGA AAAACGCCTC

2641 ACAGCTCAGC CCTACCGAGA CGGCACGCAT CATTGCCTGC CTGCCCAATC CCAAAAAATA CAATNTAAAC CCGCCAAGTG

2721 CCTACATCTC AAAACGCGGA CAATGGATTC TGCGCCAAGT GCGAACTTG AAAGGCGATA GGGCTCTGAG CGAGATTGTG

2801 AACACGCCCT AACGCCTGCC TCAACTCTTT GCACACAGTT TACCAACTCT CTGCGAAGAG TTCACAACT CTTCGCACAC

2881 ACTTCCCAA GTCTTTGCAA AGAGTTGGGA GATACTTAGG CACAAAAAAA AGGAACCTCA TGAATAGAGG TTCCCTCTTC

2961 CTTAAAAGGA ATAAATAATA ATGTTTTTTA AGCTTTAGGC TTGGCTACTT TTTCAAAGCC TGCTGCCTTC ATGCTATCTA

-----  
HindIII

3041 GGATACGCTT GCCTGGGCGG TAGTTTACGC CTACCTTTTT GATTAAGCCC GAATGAAAAT CTTTCTCTGT ATCTGCCGCT  
<<.....R8.....<

3121 CCACTGCTTA AAGTGGCATA GAGCGAGCCA AGCTTATCTA AACGAACGAT TTTGCCCGCT GCCAAGGCGT CTTGAATTAC  
<R8.<<AAGCTTAAG

-----  
HindIII

-----  
HindIII

3201 ATTCTCTAGC GCAATGATAA CGCCACGAAT ATCTGCCTCG CTGAGTGCCG AAAACTTCTC GATTGCTTA ACGAGCTGGT

3281 CTATATCCAT TTCTCCATCG CTGCCCACCA CGGCATAGTA TTTTGTGGC TCCCCTGGCT TGCTTGGGT TCTACGCTGA

3361 ATTACATTGT ATTTTATGCT CATAATTACT CTATTTTTAA TAGCCTCCCG ATGGATATAA AGTTACGCTA CAATTAGGGT

3441 CTCCATAAGC AAATCTATAC CCCTCTCTTT CATATCCCT TCTCATCTT CTTGCTCCAT CTCTCAAGGC ATCCGCTCTA

3521 TTAAGTCTAT ACCCTCCTG AAGAAATGTG TCTGCACTTG AAGAAGAATA TGAAGAGCTA TGAGAATCGT-GCAACATAGT

3601 CCAAGCTCCA TCTTGAGCTA TAACATTTGC ATGACATGTA ACACCTATAG TATAATAAAA TCTCCTAGGA GGTGTGTTC

3681 CACCACCACC TCCAGAGCTA CTACTTTTTT TACATTGTCC ATTTTGGTTA GCATGATTTT GTCCGCCATC ACTTACTAAC

3761 TTCTTAGCTT CTGCTAAGGC TTTTCTCTT GCTTCTTTT CAGCATCTGC TTGGCTAATT CCACTCACTG CTGTAGCTGT

3841 CGCTTCTTTT TTATAGTTTA CCGAGGTTC ATAATAGCCA CTACTACAA TGTTCCTGT AAAGTTTTTA TTAAGATT

3921 GAGTTTGTGT TGAGGTGTAC CCTCCGAAAC CTTTACTTC TACAGTAAAG GTAGAACTCC CCATGCTTAC GGGGAAGGTG

4001 GCGATAGTAT ACGATTGCCC TGCCGGCATT TGTTTTACTT GATACTCTC ATCTCCTCCC ACTTCTATGC TTGCCGTAA

---

4081 ATTACCACTA CCGCTAAAAG AGCCTTCTGC TATTTT TAGT GTTAAATCAT TTATATCCCC TCCTTGCCT TTTGCAGAAG  
4161 CTTTGTAC ACTTACAGCA TCATAAGCTC CTTTCCATT GGTATAAGGT ATTTATATGG CCAAAC

---

Figure 2B.

```

1  TAAAGCTGTA AWTGCTATA AACGCCCTTT AGGATAAAAT CTGCCATTTT TTGCAGTATT TTWATAGCTA AAATTTAGAA
   >>.....FrecAOR1.....>>

81  AACACCATCT CGAGTAAAGG AGCGTGTAGT GCTCGCCATC GTTGAGCGAT TGCCCACCCT CAATTGATT GGGCGAATAC
   CTTAAGCTT>>.....F6.....>>
   -----
   HindIII

161  TTGAAATAAA TGGCATCTTC TAGCGACACA TTTTGCAGCAG AAATCATGCA AAAAGCCCCG CATAAAAAGC TGAATAAAAA
241  WGCTAWTYTT CTGTTTTAAA AAAACTCATA AATTCCCCCA AATATAGAAA TATTCTGTGA AAAGTTGCAA TTTATTAACA
   <<....<

321  CTATGTGCTT GCTTTTAATG AAAAAAGTAG ATTATTTTTC CGAATCCGAA AGTTTATTTA CGCCCCATCC GATGCCTAGT
   <..FrecA-4...<<

401  CCCMSCGATA GCCATGATTA ATACAATAC AATTAAATCA WATTTTTCMC MTWWACCATA GCACAACACT TGCTAGCTCA
481  ACGAGTACTA GAGTGGTAAA AAGGATTTTT TGACGATTAT TCATGATTTT ATTTTCTCA AAGGTAAATA TTTTAAACCA
561  TAATTTTACA AATCTTAAAA TCTATTTAAA TAATAGAGAA ACCAGAAAAA AATCGTATTT TTACGGAATG AATAAAATGT
641  TACAAGTAGG CGATAAAATG CCCGATTTC AAGGTGTAGA CCAATTTGGG AAGGAGCATT CATCTGCCGA TTCAAAAAAT
721  CAGAAATTAG TCGTTTTTTT CTACCCAAAA GCCAGTACGC CAGGTTGCAC GGCAGAGGCT TGCAACATCA ACGATAATCT
801  TGATGCGCTA AAAGCACAAAG GCTACCAAGT GATAGGCGTG AGTGCAGATT CGGTAGAAAA ACAACGAAAA TTCAGTGATA
881  AATACGATTT TAAATTCCTT GTGATTGCCG ATGTGGATAA GAAAATTATT GAAGCATTTG GCGTGTGGGG CGAAAAGAAA
961  TTCATGGGTA AAACCTATGA CGGAATTCAT CGTACGACAT TCATTATTGA TGAAAACGGA GTGGTGGAGC GCGTGATAGA
   >>.....F7.....>>
   -----
   EcoRI

1041  AAAAGTGAAA ACAAAGATC ATACCAATCA AATTTTAAAT TCAGAAAAAT AAAAATATGA GCGAAATAGA CGAAGCGAAA
   >>.....recA.....>

1121  AGGAAAGCAC TCCAGCTAGT GCTTGATAAA ATGGACAAA GCTATGGTAA AGGTGCCGTG ATGATGATGG GCGACAAAGC
   >.....recA.....>
   <<.....OER1.....<

1201  CATAGACGAA AATATTCCAG TAATCCCTAC GGGGTCTCTA GGTTTAGATT TAGCCTTGGG CGTGGGAGGG TATCCGCGCG
   >.....recA.....>
   <CGAGATCTCGTGCGTGGGT
   -----
   BglIII

1281  GTAGAATCGT GGAGATTAC GGTCCAGAAT CTCTGGTAA AACCACCTTG GCAATTCATG CCATTGCCGA AGCTCAAAAG
   >.....recA.....>

1361  TCTGGCGGAA TTGCAGCTTT CATCGATGCA GAGCAGCAT TTGATAGATA TTACGCAGAA AAATTAGGCG TAGATGTTGA
   >.....recA.....>

1441  GCATTTAATT ATCTCTCAGC CAGATAATGG GGAGCAAGCT TTAGAAATTG CCGATAACTT AATCCGTTC GGTGCAATTG
   >.....recA.....>
   -----
   HindIII

1521  ATATTATTGT AATCGATTCT GTAGCGGCTT TAACGCCAAA GTCGGAAATC GACGGAGATA TGGGCGATTC CAAAATGGGA
   >.....recA.....>

1601  TTGCAAGCGC GTTGATGTC TCAAGCCTTG AGAAAGCTCA CGGGAATAT CAATAAAACC AAATGTACTG CTATTTTCAT
   >.....recA.....>

```

1681 CAACCAATTG AGAGAGAAAA TCGGTGTGAT GTTCGGTAGT CCAGAAACCA CAACGGGTGG TAATGCACTT AAATTCTATG  
 >.....recA.....>  
 1761 CATCGGTGCG TCTAGACATT CGTCGTTCTA CTCAGATTAA AGATGGGAAC GATGTCATCG GAAACTTGAC TCGCGTAAAA  
 >.....recA.....>  
 1841 GTAGTGAAAA ACAAAGTAGC TCCGCCATTC CGTAGTGCAG AATTCGACAT TATGTATGGC GAAGGAATCT CTAAAGCAGG  
 >.....recA.....>  
 -----  
 EcoRI  
 1921 CGAGATTTTA GACATTGCTA CCGATTAGA AATCGTGAAA AAAAGTGGCT CTGGGTATTC TTATGCAGAT ACTAAACTAG  
 >.....recA.....>  
 2001 GACAAGGGCG AGATGCCGTG CGTGC GTAT TGAAAGATAA TCCAGAATTA GCCGAAGAAT TAGAAGAGAA AATTAAAGAA  
 >.....recA.....>  
 CGAGATCT>>.....OEF1.....>>  
 -----  
 BglIII  
 2081 GAATTAGAGA AAAAATAGAT TTTTAGTTT TTTTAATTAA ACGAAAAATC CGTTCACTTT GTTGAACGGA TTTTTTTATG  
 >.....recA.....>  
 2161 CTTGAATGAA TTTATTTCCA ATGGATTGAA TAGCCATGCA CTTTAAATC TTCGCTATCA TAAGTGATTT CTTTGTCCGT  
 2241 GTTGGGATAG CAAACTTTAA GTCCTGCGTA TTTGGCAATG GCATGTCCTG CGGCAATGTC CAAAAGTTT ACAGGTCTAA  
 2321 AGCGGGTGTA CTCCGTAGCC CACCGATCGG CAATTAGCCC AAGTTTGATA ACGCTTCCCA TAGGCTTTGT GCGGAAAATT  
 2401 TCATGTTTCG ATTTAATTTT TTTGATGTAT TCCTCGGTGC CAGGATCCAT GTGGAATTTG CTACAAAGAA AAGTGTAATC  
 2481 TTCGGGCAAA TCCATGGTAG GAATTGGCTT GCTGTGTTT ATCAATTGTT CAAAAAATC CGATTTTCAGA GCCATTTTGT  
 2561 GCAATTGTTG TTGAGTCCCG ATGAATTTAC GAGAAGGGCA TTTATCGCTA CCGAAATAGA ACAATCCAAG CGATGGGGCG  
 2641 TACAAAATC CTAGCTTAGC CGTATTATTC TCAACTAAGC CTAGACACAC GCAATATTCA TCTGTTTTGT TGACAAAATC  
 2721 CATGGTGCCA TCAATAGGGT CTGCAATCCA ATAGGTGGGC GTATTCTTAA TTTCTTGTA AGAATCCTTA TCTCCTTCCT  
 2801 CACTAAAGTA TGGAATGTCT GTAAAGGAAA CATGTTTTTG CAAGATTTTG TTGGCGGCTA AATCTGCACT TGTAACAGGC  
 2881 GATCCGTCGG CTTTGGTCTC GGTGGAGAAT CCGTTTTGGA TTGTTTTAAA ACCTCTTCGC CAGCAAGTGC TACAGCCCGT  
 2961 GTTGCATTT CTAATAAATT CATAATCATT CTTTATTCT CGAACAAAGT CAAATAATC TCTGTATTAA AAAATAATTT  
 3041 TGGCGATAAA AATTAAATTT TATATATAAA ATATCTCTGC AAAAAACCA ATCAAATATT TAGTGAAATA AAAAAATTA  
 3121 GATTGTAAAT TTGCCTTATG TTTTAGAGA ATACCATAAA TCATAGAAAA AATACGGGCT GGATCGAAGT AATCTGTGGC  
 3201 TCTATGTTTT CGGGCAAAAC CGAAGAGTTG ATTCGTAGAG TGAAACGAGC CGAATTGGCT GGGCAAAAGG TAGAAATCTT  
 <<.....R5.....<<AAGCTTAAG  
 -----  
 HindIII  
 3281 TAAACCCGCA ATTGATAAAC GCTACGATGA GCAAGATGTG GTATCGCATG ATGAAAACAA AAAACAAGCA ACCCCGATTG  
 3361 AGGCGAGTTC TAAGTTGCCC ATTTTAGCAA GCGATTGTGA TGTGGTGGGG ATAGATGAGG CTCAATCTT TGACGAAGGA  
 3441 ATTGTTGAGG TGGCAATCT TTTAGCTAAT TCGGGGAAAA GAATAATTAT TGCGGGATTA GACATGGATT TTAAAGGTG  
 <<.....RrecAORI.....<<  
 3521 TCCATTTGGT CCTATGCCAA ATTTAATGGC GGTAGCGGAA TATGTGACCA AAGTGCATGC AATCTGTGTG AAAACAGGGA